
## NO CALCULATOR IS ALLOWED FOR THESE QUESTIONS.



The function f is differentiable on the closed interval [-6, 5] and satisfies f(-2) = 7. The graph of f', the derivative of f, consists of a semicircle and three line segments, as shown in the figure above.

- (a) Find the values of f(-6) and f(5).
- (b) On what intervals is f increasing? Justify your answer.
- (c) Find the absolute minimum value of f on the closed interval [-6, 5]. Justify your answer.
- (d) For each of f''(-5) and f''(3), find the value or explain why it does not exist.

## ANSWER WITH GRADING RUBRIC:

(a) 
$$f(-6) = f(-2) + \int_{-2}^{-6} f'(x) dx = 7 - \int_{-6}^{-2} f'(x) dx = 7 - 4 = 3$$
  
 $f(5) = f(-2) + \int_{-2}^{5} f'(x) dx = 7 - 2\pi + 3 = 10 - 2\pi$ 

- 3:  $\begin{cases} 1 : \text{ uses initial condition} \\ 1 : f(-6) \\ 1 : f(5) \end{cases}$
- (b) f'(x) > 0 on the intervals [-6, -2) and (2, 5). Therefore, f is increasing on the intervals [-6, -2] and [2, 5].
- 2 : answer with justification
- (c) The absolute minimum will occur at a critical point where f'(x) = 0 or at an endpoint.

2: 
$$\begin{cases} 1 : considers \ x = 2 \\ 1 : answer with justification \end{cases}$$

$$f'(x) = 0 \Rightarrow x = -2, x = 2$$

$$\begin{array}{c|cc}
x & f(x) \\
-6 & 3 \\
-2 & 7 \\
2 & 7 - 2\pi \\
5 & 10 - 2\pi
\end{array}$$

The absolute minimum value is  $f(2) = 7 - 2\pi$ .

(d) 
$$f''(-5) = \frac{2-0}{-6-(-2)} = -\frac{1}{2}$$

$$\lim_{x \to 3^{-}} \frac{f'(x) - f'(3)}{x - 3} = 2 \text{ and } \lim_{x \to 3^{+}} \frac{f'(x) - f'(3)}{x - 3} = -1$$


f''(3) does not exist because

$$\lim_{x \to 3^{-}} \frac{f'(x) - f'(3)}{x - 3} \neq \lim_{x \to 3^{+}} \frac{f'(x) - f'(3)}{x - 3}.$$

2:  $\begin{cases} 1: f''(-5) \\ 1: f''(3) \text{ does not exist,} \\ \text{with explanation} \end{cases}$ 

## **ANSWER WITH GRADING RUBRIC:**

No calculator is allowed for these problems.



Graph of f

The figure above shows the graph of the piecewise-linear function f. For  $-4 \le x \le 12$ , the function g is defined by  $g(x) = \int_2^x f(t) dt$ .

- (a) Does g have a relative minimum, a relative maximum, or neither at x = 10? Justify your answer.
- (b) Does the graph of g have a point of inflection at x = 4? Justify your answer.
- (c) Find the absolute minimum value and the absolute maximum value of g on the interval  $-4 \le x \le 12$ . Justify your answers.
- (d) For  $-4 \le x \le 12$ , find all intervals for which  $g(x) \le 0$ .

(a) The function 
$$g$$
 has neither a relative minimum nor a relative maximum at  $x = 10$  since  $g'(x) = f(x)$  and  $f(x) \le 0$  for  $8 \le x \le 12$ .

- (b) The graph of g has a point of inflection at x = 4 since g'(x) = f(x) is increasing for  $2 \le x \le 4$  and decreasing for  $4 \le x \le 8$ .
- (c) g'(x) = f(x) changes sign only at x = -2 and x = 6.

$$\begin{array}{c|cc}
x & g(x) \\
-4 & -4 \\
-2 & -8 \\
6 & 8 \\
12 & -4
\end{array}$$

On the interval  $-4 \le x \le 12$ , the absolute minimum value is g(-2) = -8 and the absolute maximum value is g(6) = 8.

(d) 
$$g(x) \le 0$$
 for  $-4 \le x \le 2$  and  $10 \le x \le 12$ .

1: 
$$g'(x) = f(x)$$
 in (a), (b), (c), or (d)

1: answer with justification


1: answer with justification

1: considers 
$$x = -2$$
 and  $x = 6$   
as candidates

1: considers x = -4 and x = 12

2: answers with justification

2: intervals



Graph of  $\nu$ 

A particle moves along the x-axis so that its velocity at time t, for  $0 \le t \le 6$ , is given by a differentiable function  $\nu$  whose graph is shown above. The velocity is 0 at t = 0, t = 3, and t = 5, and the graph has horizontal tangents at t = 1 and t = 4. The areas of the regions bounded by the t-axis and the graph of  $\nu$  on the intervals [0, 3], [3, 5], and [5, 6] are [5