

AP Calculus AB:
Section I, Part A

50 Minutes—No Calculator

Note: Unless otherwise specified, the domain of a function f is assumed to be the set of all real numbers x for which $f(x)$ is a real number.

1. $\int_1^2 (4x^3 - 6x) dx =$

$$x^4 - 3x^2 \Big|_1^2$$

(A) 2

(B) 4

(C) 6

(D) 36

(E) 42

$$(16 - 12) - (1 - 3)$$

$$4 - (-2) = 6$$

2. If $f(x) = x\sqrt{2x-3}$, then $f'(x) =$

(A) $\frac{3x-3}{\sqrt{2x-3}}$

(B) $\frac{x}{\sqrt{2x-3}}$

(C) $\frac{1}{\sqrt{2x-3}}$

(D) $\frac{-x+3}{\sqrt{2x-3}}$

(E) $\frac{5x-6}{2\sqrt{2x-3}}$

$$f'(x) = x \left(\frac{1}{2} (2x-3)^{-1/2} \cdot 2 \right) + \sqrt{2x-3}$$

$$= \frac{x}{\sqrt{2x-3}} + \frac{2x-3}{\sqrt{2x-3}}$$

$$= \frac{3x-3}{\sqrt{2x-3}}$$

3. If $\int_a^b f(x) dx = a + 2b$, then $\int_a^b (f(x) + 5) dx =$

(A) $a + 2b + 5$

(B) $5b - 5a$

(C) $7b - 4a$

(D) $7b - 5a$

(E) $7b - 6a$

$$\int_a^b f(x) dx + \int_a^b 5 dx$$

$$a + 2b + 5b - 5a = -4a + 7b$$

4. If $f(x) = -x^3 + x + \frac{1}{x}$, then $f'(-1) =$

(A) 3

(B) 1

(C) -1

(D) -3

(E) -5

$$f'(-1) = -3(-1)^2 + 1 - \frac{1}{(-1)^2} = -3 + 1 - 1 = -3$$

5. The graph of $y = 3x^4 - 16x^3 + 24x^2 + 48$ is concave down for

- (A) $x < 0$
 (B) $x > 0$
 (C) $x < -2$ or $x > -\frac{2}{3}$
 (D) $x < \frac{2}{3}$ or $x > 2$
 (E) $\frac{2}{3} < x < 2$

$$y' = 12x^3 - 48x^2 + 48x$$

$$y'' = 36x^2 - 96x + 48$$

$$0 = 12(3x^2 - 8x + 4)$$

$$= 12(3x - 2)(x - 2)$$

-	-	+	-	+	+
P	2/3	N	2	P	P

6. $\frac{1}{2} \int e^{\frac{t}{2}} dt =$

- (A) $e^{-1} + C$ (B) $e^{-\frac{1}{2}} + C$ (C) $e^{\frac{1}{2}} + C$ (D) $2e^{\frac{1}{2}} + C$ (E) $e^1 + C$

$$u = \frac{1}{2}t \quad du = \frac{1}{2}dt \quad \int e^u du = e^u + C$$

7. $\frac{d}{dx} \cos^2(x^3) =$


- (A) $6x^2 \sin(x^3) \cos(x^3)$
 (B) $6x^2 \cos(x^3)$
 (C) $\sin^2(x^3)$
 (D) $-6x^2 \sin(x^3) \cos(x^3)$
 (E) $-2 \sin(x^3) \cos(x^3)$

$$\frac{d}{dx} (\cos(x^3))^2$$

$$= 2(\cos x^3) \cdot (-\sin x^3) \cdot 3x^2$$

$$= -6x^2 \cos(x^3) \sin(x^3)$$

Questions 8-9 refer to the following situation.



$v(t)$ changes signs

A bug begins to crawl up a vertical wire at time $t = 0$. The velocity v of the bug at time t , $0 \leq t \leq 8$, is given by the function whose graph is shown above.

8. At what value of t does the bug change direction?
 (A) 2 (B) 4 (C) 6 (D) 7 (E) 8

9. What is the total distance the bug traveled from $t = 0$ to $t = 8$?
 (A) 14 (B) 13 (C) 11 (D) 8 (E) 6

10. An equation of the line tangent to the graph of $y = \cos(2x)$ at $x = \frac{\pi}{4}$ is


- (A) $y - 1 = -\left(x - \frac{\pi}{4}\right)$
 (B) $y - 1 = -2\left(x - \frac{\pi}{4}\right)$
 (C) $y = 2\left(x - \frac{\pi}{4}\right)$
 (D) $y = -\left(x - \frac{\pi}{4}\right)$
 (E) $y = -2\left(x - \frac{\pi}{4}\right)$

$$y' = -\sin(2x) \cdot 2$$






$$= -2\sin\left(2\left(\frac{\pi}{4}\right)\right)$$

$$= -2\sin\frac{\pi}{2} = -2 \quad m = -2$$

$$y = \cos\frac{\pi}{2} = 0 \quad \left(\frac{\pi}{4}, 0\right)$$



11. The graph of the derivative of f is shown in the figure above. Which of the following could be the graph of f ?



12. At what point on the graph of $y = \frac{1}{2}x^2$ is the tangent line parallel to the line $2x - 4y = 3$?

(A) $(\frac{1}{2}, -\frac{1}{2})$

(B) $(\frac{1}{2}, \frac{1}{8})$

(C) $(1, -\frac{1}{4})$

(D) $(1, \frac{1}{2})$

(E) $(2, 2)$

$y' = x$

$\frac{1}{2} = x$

$y = \frac{1}{2}(\frac{1}{2})^2 = \frac{1}{2}(\frac{1}{4}) = \frac{1}{8}$


$-4y = -2x + 3$

$y = \frac{1}{2}x - \frac{3}{4}$

AP Calculus AB:
Section I, Part A

13. Let f be a function defined for all real numbers x . If $f'(x) = \frac{4-x^2}{x-2}$, then f is decreasing on the interval
- when $x-2 < 0$
 $x < 2$*
- (A) $(-\infty, 2)$ (B) $(-\infty, \infty)$ (C) $(-2, 4)$ (D) $(-2, \infty)$ (E) $(2, \infty)$

14. Let f be a differentiable function such that $f(3) = 2$ and $f'(3) = 5$. If the tangent line to the graph of f at $x = 3$ is used to find an approximation to a zero of f , that approximation is
- (A) 0.4 (B) 0.5 (C) 2.6 (D) 3.4 (E) 5.5
- $y - 2 = 5(x - 3)$
 $0 - 2 = 5x - 15$
 $13 = 5x$*




15. The graph of the function f is shown in the figure above. Which of the following statements about f is true?

- (A) $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow b} f(x)$
- (B) $\lim_{x \rightarrow a} f(x) = 2$
- (C) $\lim_{x \rightarrow b} f(x) = 2$
- (D) $\lim_{x \rightarrow b} f(x) = 1$
- (E) $\lim_{x \rightarrow a} f(x)$ does not exist.

AP Calculus AB:
Section I, Part A

16. The area of the region enclosed by the graph of $y = x^2 + 1$ and the line $y = 5$ is

- (A) $\frac{14}{3}$ (B) $\frac{16}{3}$ (C) $\frac{28}{3}$ (D) $\frac{32}{3}$ (E) 8π



17. If $x^2 + y^2 = 25$, what is the value of $\frac{d^2y}{dx^2}$ at the point $(4, 3)$?

- (A) $-\frac{25}{27}$ (B) $-\frac{7}{27}$ (C) $\frac{7}{27}$ (D) $\frac{3}{4}$ (E) $\frac{25}{27}$

18. $\int_0^{\pi/4} \frac{e^{\tan x}}{\cos^2 x} dx$ is $\int_0^{\pi/4} \sec^2 x e^{\tan x} dx$

- (A) 0 (B) 1 (C) $e - 1$ (D) e (E) $e + 1$

$u = \tan x$ $u(\pi/4) = 1$
 $du = \sec^2 x dx$ $u(0) = 0$
 $\int_0^1 e^u du = e^1 - e^0 = e - 1$

19. If $f(x) = \ln|x^2 - 1|$, then $f'(x) = \frac{1}{x^2 - 1} (2x) = \frac{2x}{x^2 - 1}$

- (A) $\left| \frac{2x}{x^2 - 1} \right|$
 (B) $\frac{2x}{x^2 - 1}$
 (C) $\frac{2|x|}{x^2 - 1}$
 (D) $\frac{2x}{x^2 - 1}$
 (E) $\frac{1}{x^2 - 1}$

$$17. x^2 + y^2 = 25$$

$$2x + 2y y' = 0$$

$$2y y' = -2x$$

$$y' = -\frac{2x}{2y}$$

$$y' = -\frac{x}{y}$$

$$y'' = \frac{y(-1) - (-x)y'}{y^2}$$

$$y'' = \frac{-y + x\left(-\frac{x}{y}\right)}{y^2}$$

$$= \frac{-y + \frac{-x^2}{y}}{y^2} =$$

$$= \frac{\frac{-y^2 - x^2}{y}}{y^2} = \frac{-1(x^2 + y^2)}{y^3} = \frac{-25}{y^3} = \frac{-25}{(3)^3} = \frac{-25}{27}$$

$$\sin(-x) = -\sin x$$

AP Calculus AB:
Section I, Part A

20. The average value of $\cos x$ on the interval $[-3, 5]$ is

(A) $\frac{\sin 5 - \sin 3}{8}$

(B) $\frac{\sin 5 - \sin 3}{2}$

(C) $\frac{\sin 3 - \sin 5}{2}$

(D) $\frac{\sin 3 + \sin 5}{2}$

(E) $\frac{\sin 3 + \sin 5}{8}$

$$\begin{aligned} \frac{1}{5 - (-3)} \int_{-3}^5 \cos x \, dx &= \frac{1}{8} (\sin x) \Big|_{-3}^5 \\ &= \frac{\sin 5 - (\sin(-3))}{8} \\ &= \frac{\sin 5 + \sin 3}{8} \end{aligned}$$

odd function

21. $\lim_{x \rightarrow 1} \frac{x}{\ln x}$ is

(A) 0

(B) $\frac{1}{e}$

(C) 1

(D) e

(E) nonexistent

22. What are all values of x for which the function f defined by $f(x) = (x^2 - 3)e^{-x}$ is increasing?

(A) There are no such values of x .

(B) $x < -1$ and $x > 3$

(C) $-3 < x < 1$

(D) $-1 < x < 3$

(E) All values of x

$$\begin{aligned} f'(x) &= (x^2 - 3)(-e^{-x}) + e^{-x}(2x) \\ &= e^{-x}(-x^2 + 3 + 2x) \\ &= -e^{-x}(x^2 - 2x - 3) \\ &= -e^{-x}(x-3)(x+1) \end{aligned}$$

$\begin{matrix} + & - & - & + \\ \hline N & -1 & P & 3 & N \end{matrix}$

23. If the region enclosed by the y -axis, the line $y = 2$, and the curve $y = \sqrt{x}$ is revolved about the y -axis, the volume of the solid generated is

(A) $\frac{32\pi}{5}$

(B) $\frac{16\pi}{3}$


(C) $\frac{16\pi}{5}$

(D) $\frac{8\pi}{3}$

(E) π

$$\int_0^2 \pi y^4 \, dy$$

$$\frac{5}{5} \pi \Big|_0^2 = \frac{32\pi}{5}$$



24. The expression $\frac{1}{50} \left(\sqrt{\frac{1}{50}} + \sqrt{\frac{2}{50}} + \sqrt{\frac{3}{50}} + \dots + \sqrt{\frac{50}{50}} \right)$ is a Riemann sum approximation for

interval 0, 1
 $f(x) = \sqrt{x}$

$\frac{1-0}{50} = \frac{1}{50}$


(A) $\int_0^1 \sqrt{\frac{x}{50}} dx$

(B) $\int_0^1 \sqrt{x} dx$

(C) $\frac{1}{50} \int_0^1 \sqrt{\frac{x}{50}} dx$

(D) $\frac{1}{50} \int_0^1 \sqrt{x} dx$

(E) $\frac{1}{50} \int_0^{50} \sqrt{x} dx$



25. $\int x \sin(2x) dx =$

(A) $-\frac{x}{2} \cos(2x) + \frac{1}{4} \sin(2x) + C$

(B) $-\frac{x}{2} \cos(2x) - \frac{1}{4} \sin(2x) + C$

(C) $\frac{x}{2} \cos(2x) - \frac{1}{4} \sin(2x) + C$

(D) $-2x \cos(2x) + \sin(2x) + C$

(E) $-2x \cos(2x) - 4 \sin(2x) + C$

$x \cdot 2 \sin x \cos x$

$2 \times \sin x \cos x$

Don't assign
be parts
do