

**AP<sup>®</sup> CALCULUS AB**  
**2007 SCORING GUIDELINES**

**Question 5**

$t$ (minutes)	0	2	5	7	11	12
$r'(t)$ (feet per minute)	5.7	4.0	2.0	1.2	0.6	0.5

The volume of a spherical hot air balloon expands as the air inside the balloon is heated. The radius of the balloon, in feet, is modeled by a twice-differentiable function  $r$  of time  $t$ , where  $t$  is measured in minutes. For  $0 < t < 12$ , the graph of  $r$  is concave down. The table above gives selected values of the rate of change,  $r'(t)$ , of the radius of the balloon over the time interval  $0 \leq t \leq 12$ . The radius of the balloon is 30 feet when  $t = 5$ . (Note: The volume of a sphere of radius  $r$  is given by  $V = \frac{4}{3}\pi r^3$ .)

- (a) Estimate the radius of the balloon when  $t = 5.4$  using the tangent line approximation at  $t = 5$ . Is your estimate greater than or less than the true value? Give a reason for your answer.
- (b) Find the rate of change of the volume of the balloon with respect to time when  $t = 5$ . Indicate units of measure.
- (c) Use a right Riemann sum with the five subintervals indicated by the data in the table to approximate  $\int_0^{12} r'(t) dt$ . Using correct units, explain the meaning of  $\int_0^{12} r'(t) dt$  in terms of the radius of the balloon.
- (d) Is your approximation in part (c) greater than or less than  $\int_0^{12} r'(t) dt$ ? Give a reason for your answer.

<p>(a) <math>r(5.4) = r(5) + r'(5)\Delta t = 30 + 2(0.4) = 30.8</math> ft Since the graph of <math>r</math> is concave down on the interval <math>5 &lt; t &lt; 5.4</math>, this estimate is greater than <math>r(5.4)</math>.</p>	<p>2 : <math>\left\{ \begin{array}{l} 1 : \text{estimate} \\ 1 : \text{conclusion with reason} \end{array} \right.</math></p>
<p>(b) <math>\frac{dV}{dt} = 3\left(\frac{4}{3}\right)\pi r^2 \frac{dr}{dt}</math> <math>\left. \frac{dV}{dt} \right _{t=5} = 4\pi(30)^2 2 = 7200\pi</math> ft<sup>3</sup>/min</p>	<p>3 : <math>\left\{ \begin{array}{l} 2 : \frac{dV}{dt} \\ 1 : \text{answer} \end{array} \right.</math></p>
<p>(c) <math>\int_0^{12} r'(t) dt \approx 2(4.0) + 3(2.0) + 2(1.2) + 4(0.6) + 1(0.5)</math> <math>= 19.3</math> ft <math>\int_0^{12} r'(t) dt</math> is the change in the radius, in feet, from <math>t = 0</math> to <math>t = 12</math> minutes.</p>	<p>2 : <math>\left\{ \begin{array}{l} 1 : \text{approximation} \\ 1 : \text{explanation} \end{array} \right.</math></p>
<p>(d) Since <math>r</math> is concave down, <math>r'</math> is decreasing on <math>0 &lt; t &lt; 12</math>. Therefore, this approximation, 19.3 ft, is less than <math>\int_0^{12} r'(t) dt</math>.</p> <p>Units of ft<sup>3</sup>/min in part (b) and ft in part (c)</p>	<p>1 : conclusion with reason</p> <p>1 : units in (b) and (c)</p>